

Status: Currently Official on 17-Feb-2025  
 Official Date: Official as of 01-Jan-2018  
 Document Type: USP Monographs  
 DocId: GUID-15869C35-50B9-4E51-A348-8FE64C493BC5\_3\_en-US  
 DOI: [https://doi.org/10.31003/USPNF\\_M2358\\_03\\_01](https://doi.org/10.31003/USPNF_M2358_03_01)  
 DOI Ref: ec5kp

© 2025 USPC  
 Do not distribute

## Aluminum Zirconium Octachlorohydrate Gly

» Aluminum Zirconium Octachlorohydrate Gly is a derivative of Aluminum Zirconium Octachlorohydrate in which some of the water molecules have been displaced by glycine, calcium glycinate, magnesium glycinate, potassium glycinate, sodium glycinate, or zinc glycinate. It encompasses a range of aluminum-to-zirconium atomic ratios between 6.0:1 and 10.0:1, and a range of (aluminum plus zirconium)-to-chloride atomic ratios between 1.5:1 and 0.9:1. It contains not less than 90.0 percent and not more than 110.0 percent of the labeled amount of anhydrous aluminum zirconium octachlorohydrate.

**Packaging and storage**—Preserve in well-closed containers.

**Labeling**—The label states the form of glycine used and the claimed content of anhydrous aluminum zirconium octachlorohydrate.

**Identification**—

**A:** A solution (1 in 10) responds to the test for [Chloride \(191\)](#).

**B:** Place about 0.5 g of it in a 50-mL beaker, add about 20 mL of water, and swirl to dissolve. Heat to boiling on a hot plate, and add about 60 mg of ninhydrin: a deep violet color immediately develops.

**pH (791):** between 3.0 and 5.0, in a solution [15 in 100 (w/w)].

**ARSENIC, Method I (211):** 2 µg per g.

**Limit of iron**—Using Aluminum Zirconium Octachlorohydrate Gly instead of Aluminum Zirconium Octachlorohydrate, proceed as directed in the test for *Limit of iron* under [Aluminum Zirconium Octachlorohydrate](#). The specified result is obtained (150 µg per g limit).

**Content of aluminum**—Using Aluminum Zirconium Octachlorohydrate Gly instead of Aluminum Zirconium Octachlorohydrate, proceed as directed in the test for *Content of aluminum* under [Aluminum Zirconium Octachlorohydrate](#). Use the result obtained to calculate the *Aluminum/zirconium atomic ratio* and the *(Aluminum plus zirconium)/chloride atomic ratio*.

**Content of zirconium**—Using Aluminum Zirconium Octachlorohydrate Gly instead of Aluminum Zirconium Octachlorohydrate, proceed as directed in the test for *Content of zirconium* under [Aluminum Zirconium Octachlorohydrate](#). Use the result obtained to calculate the *Aluminum/zirconium atomic ratio* and the *(Aluminum plus zirconium)/chloride atomic ratio*.

**Aluminum/zirconium atomic ratio**—Divide the percentage of aluminum found in the test for *Content of aluminum* by the percentage of zirconium found in the test for *Content of zirconium*, and multiply by 92.97/26.98, in which 92.97 is the atomic weight of zirconium corrected for 2% hafnium content, and 26.98 is the atomic weight of aluminum: the ratio is between 6.0:1 and 10.0:1.

**Content of chloride**—Using Aluminum Zirconium Octachlorohydrate Gly instead of Aluminum Zirconium Octachlorohydrate, proceed as directed in the test for *Content of chloride* under [Aluminum Zirconium Octachlorohydrate](#). Use the result obtained to calculate the *(Aluminum plus zirconium)/chloride atomic ratio*.

**(Aluminum plus zirconium)/chloride atomic ratio**—Calculate the (aluminum plus zirconium)/chloride atomic ratio by the formula:

$$[(Al/26.98) + (Zr/92.97)]/(Cl/35.453)$$

in which *Al*, *Zr*, and *Cl* are the percentages of aluminum, zirconium, and chloride as determined in the tests for *Content of aluminum*, *Content of zirconium*, and *Content of chloride*, respectively; 26.98 is the atomic weight of aluminum; 92.97 is the atomic weight of zirconium corrected for 2% hafnium content; and 35.453 is the atomic weight of chlorine: the ratio is between 1.5:1 and 0.9:1.

**Assay**—Calculate the percentage of anhydrous aluminum zirconium octachlorohydrate in the Aluminum Zirconium Octachlorohydrate Gly by the formula:

$$Al\{26.98y + 92.97 + 17.01[3y + 4 - (y + 1)/z] + 35.453(y + 1)/z\}/26.98y$$

in which *Al* is the percentage of aluminum found in the test for *Content of aluminum*, *y* is the aluminum/zirconium atomic ratio found in the test for *Aluminum/zirconium atomic ratio*, *z* is the (aluminum plus zirconium)/chloride atomic ratio found in the test for *(Aluminum plus zirconium)/chloride atomic ratio*, 26.98 is the atomic weight of aluminum, 92.97 is the atomic weight of zirconium corrected for 2% hafnium content, 17.01 is the molecular weight of the hydroxide anion (OH), and 35.453 is the atomic weight of chlorine (Cl).

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

| Topic/Question                          | Contact                                       | Expert Committee          |
|-----------------------------------------|-----------------------------------------------|---------------------------|
| ALUMINUM ZIRCONIUM OCTACHLOROXYDREX GLY | <a href="#">Documentary Standards Support</a> | SM32020 Small Molecules 3 |



Documents

Document Sharing Hub

| Topic/Question             | Contact                                                                     | Expert Committee          |
|----------------------------|-----------------------------------------------------------------------------|---------------------------|
| REFERENCE STANDARD SUPPORT | RS Technical Services<br><a href="mailto:RSTECH@usp.org">RSTECH@usp.org</a> | SM32020 Small Molecules 3 |

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 27(4)

Current DocID: GUID-15869C35-50B9-4E51-A348-8FE64C493BC5\_3\_en-US

Previous DocID: GUID-15869C35-50B9-4E51-A348-8FE64C493BC5\_1\_en-US

DOI: [https://doi.org/10.31003/USPNF\\_M2358\\_03\\_01](https://doi.org/10.31003/USPNF_M2358_03_01)

DOI ref: [ec5kp](#)

OFFICIAL