

Status: Currently Official on 17-Feb-2025
 Official Date: Official as of 01-Jun-2023
 Document Type: USP Monographs
 DocId: GUID-7E81E65F-5BF2-458C-9131-542A42CC84F8_4_en-US
 DOI: https://doi.org/10.31003/USPNF_M2368_04_01
 DOI Ref: 187gq

© 2025 USPC
 Do not distribute

Aluminum Zirconium Tetrachlorohydrate Gly

» Aluminum Zirconium Tetrachlorohydrate Gly is a derivative of Aluminum Zirconium Tetrachlorohydrate in which some of the water molecules have been displaced by glycine, calcium glycinate, magnesium glycinate, potassium glycinate, sodium glycinate, or zinc glycinate. It encompasses a range of aluminum-to-zirconium atomic ratios between 2.0:1 and 5.99:1 and a range of (aluminum plus zirconium)-to-chloride atomic ratios between 1.5:1 and 0.9:1. It contains not less than 90.0 percent and not more than 110.0 percent of the labeled amount of anhydrous aluminum zirconium tetrachlorohydrate.

Packaging and storage—Preserve in well-closed containers.

Labeling—The label states the form of glycine used and the claimed content of anhydrous aluminum zirconium tetrachlorohydrate.

Identification—

A: A solution (1 in 10) responds to the test for [Chloride \(191\)](#).

B: Place about 0.5 g of it in a 50-mL beaker, add about 20 mL of water, and swirl to dissolve. Heat to boiling on a hot plate, and add about 60 mg of ninhydrin: a deep violet color immediately develops.

pH (791): between 3.0 and 5.0, in a solution [15 in 100 (w/w)].

Change to read:

▲ [ARSENIC \(211\), Procedures, Procedure 1](#) ▲ (CN 1-Jun-2023) : 2 µg per g.

Limit of iron—Using Aluminum Zirconium Tetrachlorohydrate Gly instead of Aluminum Zirconium Octachlorohydrate, proceed as directed in the test for *Limit of iron* under [Aluminum Zirconium Octachlorohydrate](#). The specified result is obtained (150 µg per g limit).

Content of aluminum—Using Aluminum Zirconium Tetrachlorohydrate Gly instead of Aluminum Zirconium Octachlorohydrate, proceed as directed in the test for *Content of aluminum* under [Aluminum Zirconium Octachlorohydrate](#). Use the result obtained to calculate the *Aluminum/zirconium atomic ratio* and the *(Aluminum plus zirconium)/chloride atomic ratio*.

Content of zirconium—Using Aluminum Zirconium Tetrachlorohydrate Gly instead of Aluminum Zirconium Octachlorohydrate, proceed as directed in the test for *Content of zirconium* under [Aluminum Zirconium Octachlorohydrate](#). Use the result obtained to calculate the *Aluminum/zirconium atomic ratio* and the *(Aluminum plus zirconium)/chloride atomic ratio*.

Aluminum/zirconium atomic ratio—Divide the percentage of aluminum found in the test for *Content of aluminum* by the percentage of zirconium found in the test for *Content of zirconium*, and multiply by 92.97/26.98, in which 92.97 is the atomic weight of zirconium corrected for 2% hafnium content, and 26.98 is the atomic weight of aluminum: the ratio is between 2.0:1 and 5.99:1.

Content of chloride—Using Aluminum Zirconium Tetrachlorohydrate Gly instead of Aluminum Zirconium Octachlorohydrate, proceed as directed in the test for *Content of chloride* under [Aluminum Zirconium Octachlorohydrate](#). Use the result obtained to calculate the *(Aluminum plus zirconium)/chloride atomic ratio*.

(Aluminum plus zirconium)/chloride atomic ratio—Calculate the (aluminum plus zirconium)/chloride atomic ratio by the formula:

$$[(Al/26.98) + (Zr/92.97)]/(Cl/35.453)$$

in which *Al*, *Zr*, and *Cl* are the percentages of aluminum, zirconium, and chloride as determined in the tests for *Content of aluminum*, *Content of zirconium*, and *Content of chloride*, respectively; 26.98 is the atomic weight of aluminum; 92.97 is the atomic weight of zirconium corrected for 2% hafnium content; and 35.453 is the atomic weight of chlorine: the ratio is between 1.5:1 and 0.9:1.

Assay—Calculate the percentage of anhydrous aluminum zirconium tetrachlorohydrate in the Aluminum Zirconium Tetrachlorohydrate Gly by the formula:

$$Al/((26.98y + 92.97 + 17.01[3y + 4 - (y + 1)/z] + 35.453(y + 1)/z)/26.98y)$$

in which *Al* is the percentage of aluminum found in the test for *Content of aluminum*, *y* is the aluminum/zirconium atomic ratio found in the test for *Aluminum/zirconium atomic ratio*, *z* is the (aluminum plus zirconium)/chloride atomic ratio found in the test for *(Aluminum plus zirconium)/chloride atomic ratio*, 26.98 is the atomic weight of aluminum, 92.97 is the atomic weight of zirconium corrected for 2% hafnium content, 17.01 is the molecular weight of the hydroxide anion (OH), and 35.453 is the atomic weight of chlorine (Cl).

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
ALUMINUM ZIRCONIUM TETRACHLOROHYDREX GLY	Documentary Standards Support	SM32020 Small Molecules 3
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	SM32020 Small Molecules 3

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 27(4)

Current DocID: GUID-7E81E65F-5BF2-458C-9131-542A42CC84F8_4_en-US

DOI: https://doi.org/10.31003/USPNF_M2368_04_01

DOI ref: [187gg](#)

OFFICIAL