

Status: Currently Official on 17-Feb-2025
Official Date: Official as of 01-Oct-2024
Document Type: USP Monographs
DocId: GUID-54821A7B-3598-496E-9DE0-0B3FB91C3483_6_en-US
DOI: https://doi.org/10.31003/USPNF_M3580_06_01
DOI Ref: ow3bg

© 2025 USPC
Do not distribute

Amlodipine and Benazepril Hydrochloride Capsules

DEFINITION

Amlodipine and Benazepril Hydrochloride Capsules contain an amount of amlodipine besylate equivalent to NLT 90.0% and NMT 110.0% of the labeled amount of amlodipine ($C_{20}H_{25}N_2O_5Cl$) and NLT 90.0% and NMT 110.0% of the labeled amount of benazepril hydrochloride ($C_{24}H_{28}N_2O_5 \cdot HCl$).

IDENTIFICATION

- A. The retention times of the major peaks of the *Sample solution* correspond to those of the *Standard solution*, as obtained in the Assay
- B. The UV spectra of the major peaks of the *Sample solution* correspond to those of the *Standard solution*, as obtained in the Assay.

ASSAY

Change to read:

PROCEDURE

Buffer 1: 0.7% (v/v) [triethylamine](#) in [water](#). Adjust with [phosphoric acid](#) to a pH of 3.0, and add 1.2 g/L of [tetrabutylammonium](#) (ERR-1-Oct-2024) [hydrogen sulfate](#) to this solution.

Buffer 2: 0.7% (v/v) [triethylamine](#) in [water](#). Adjust with [phosphoric acid](#) to a pH of 3.0.

Mobile phase: [Acetonitrile](#), [methanol](#), and *Buffer 1* (10:30:70)

Diluent: [Acetonitrile](#), [methanol](#), and *Buffer 2* (20:30:50)

Standard solution: Concentrations of [USP Amlodipine Besylate RS](#) and [USP Benazepril Hydrochloride RS](#) in *Diluent* as directed in [Table 1](#).

Table 1

Strength of Capsule Amlodipine/Benazepril Hydrochloride (mg/mg)	Concentration of USP Amlodipine Besylate RS / USP Benazepril Hydrochloride RS (mg/mL)
2.5/10	0.18/0.5
5/20	0.18/0.5
5/10	0.18/0.25
10/20	0.36/0.5
5/40 and 10/40	0.056/0.16

Sample solution: Transfer the contents of 5 Capsules into a suitable volumetric flask to obtain nominal concentrations as given in [Table 2](#). Add *Diluent* to about 70% of the volume of the flask and keep on a rotary shaker for about 45 min, sonicate for about 30 min with occasional shaking, and dilute with *Diluent* to volume. Centrifuge a portion of the solution for about 10 min, and pass through a filter of 0.45-µm pore size.

Table 2

Strength of Capsule Amlodipine/Benazepril Hydrochloride (mg/mg)	Nominal Concentration of Amlodipine/Benazepril Hydrochloride (mg/mL)
2.5/10	0.125/0.5
5/20	0.125/0.5

Strength of Capsule Amlodipine/Benazepril Hydrochloride (mg/mg)	Nominal Concentration of Amlodipine/Benazepril Hydrochloride (mg/mL)
5/10	0.125/0.25
10/20	0.25/0.5
5/40	0.02/0.16
10/40	0.04/0.16

Chromatographic system(See [Chromatography \(621\), System Suitability.](#))**Mode:** LC**Detector:** UV 237 nm. For *Identification B*, use a diode array detector in the range of 200–400 nm.**Column:** 4.6-mm × 15-cm; 5-μm packing [L1](#)**Flow rate:** 1.2 mL/min**Injection volume:** 10 μL**Run time:** NLT 2 times the retention time of amlodipine**System suitability****Sample:** *Standard solution***Suitability requirements****Tailing factor:** NMT 2.0 for the amlodipine and benazepril peaks**Relative standard deviation:** NMT 2.0% for the amlodipine and benazepril peaks**Analysis****Samples:** *Standard solution* and *Sample solution*Calculate the percentage of the labeled amount of amlodipine ($C_{20}H_{25}N_2O_5Cl$) in the portion of Capsules taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (M_{r1}/M_{r2}) \times 100$$

 r_U = peak response of amlodipine from the *Sample solution* r_S = peak response of amlodipine from the *Standard solution* C_S = concentration of [USP Amlodipine Besylate RS](#) in the *Standard solution* (mg/mL) C_U = nominal concentration of amlodipine in the *Sample solution* (mg/mL) M_{r1} = molecular weight of amlodipine, 408.88 M_{r2} = molecular weight of amlodipine besylate, 567.05Calculate the percentage of the labeled amount of benazepril hydrochloride ($C_{24}H_{28}N_2O_5 \cdot HCl$), in the portion of Capsules taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

 r_U = peak response of benazepril from the *Sample solution* r_S = peak response of benazepril from the *Standard solution* C_S = concentration of [USP Benazepril Hydrochloride RS](#) in the *Standard solution* (mg/mL) C_U = nominal concentration of benazepril hydrochloride in the *Sample solution* (mg/mL)**Acceptance criteria:** 90.0%–110.0%**PERFORMANCE TESTS**• **[DISSOLUTION \(711\)](#)****Medium:** [0.01 N hydrochloric acid](#); 500 mL**Apparatus 1:** 100 rpm**Time:** 30 min**Buffer:** 2.72 g/L of [potassium phosphate, monobasic](#) in [water](#). Add 0.2% (v/v) [triethylamine](#) per liter. Adjust with [phosphoric acid](#) to a pH of 3.0.**Mobile phase:** [Acetonitrile](#), [methanol](#), and *Buffer* (15:35:50)