

Status: Currently Official on 13-Feb-2025
Official Date: Official Prior to 2013
Document Type: USP Monographs
DocId: GUID-5BB4D572-99C6-4C70-B5F9-52106952F5C4_1_en-US
DOI: https://doi.org/10.31003/USPNF_M6885_01_01
DOI Ref: 52eqq

© 2025 USPC
Do not distribute

Bacitracin and Polymyxin B Sulfate Topical Aerosol

DEFINITION

Bacitracin and Polymyxin B Sulfate Topical Aerosol is a suspension of Bacitracin and Polymyxin B Sulfate in a suitable vehicle, packaged in a pressurized container with a suitable inert propellant. It contains NLT 90.0% and NMT 130.0% of the labeled amounts of bacitracin and polymyxin B. It may contain a suitable local anesthetic.

Prepare the specimen for the following tests and assays as follows. Maintain the container in the inverted position throughout this procedure. Store the container in a freezer at -70° for 16–24 h. Remove the container from the freezer, promptly puncture the container, and allow the propellant to volatilize. Open the container, and mix the contents.

IDENTIFICATION

• A. [THIN-LAYER CHROMATOGRAPHIC IDENTIFICATION TEST \(201BNP\)](#).

Sample: Prepare as directed above.

Analysis: Test as directed in the section *For Creams, Lotions, and Ointments* in the chapter.

Acceptance criteria: Meets the requirements

ASSAY

• BACITRACIN

(See [Antibiotics—Microbial Assays \(81\)](#).)

Sample solution: Use a portion of the contents of one container, containing nominally 500 USP Bacitracin Units, prepared as directed above. Transfer to a suitable separator containing 50 mL of ether, and extract with three 25-mL portions of *Buffer B.1* (see the chapter). Combine the buffer extracts in a 100-mL volumetric flask, dilute with *Buffer B.1* to volume, and mix.

Analysis: Proceed as directed in the chapter. Add sufficient 0.01 N hydrochloric acid to this solution so that the amount of hydrochloric acid in the *Test Dilution* is the same as in the median level of the standard. Dilute with *Buffer B.1* to obtain a *Test Dilution* having a bacitracin concentration that is nominally equivalent to the median level of the standard.

Acceptance criteria: 90.0%–130.0%

• POLYMYXIN B

(See [Antibiotics—Microbial Assays \(81\)](#).)

Sample solution: Use a portion of the contents of one container, containing nominally 5000 USP Polymyxin B Units, prepared as directed above. Transfer to a suitable separator containing 50 mL of ether, and extract with three 25-mL portions of *Buffer B.6* (see the chapter). Combine the buffer extracts in a 100-mL volumetric flask, dilute with *Buffer B.6* to volume, and mix.

Analysis: Proceed as directed in the chapter. Dilute a suitable aliquot of the *Sample solution* with *Buffer B.6* to obtain a *Test Dilution* having a polymyxin B concentration that is nominally equivalent to the median level of the standard.

Acceptance criteria: 90.0%–130.0%

SPECIFIC TESTS

• WATER DETERMINATION, [Method I \(921\)](#).

Analysis: Use a portion of the contents of one container, prepared as directed above, and 20 mL of a mixture of toluene and methanol (7:3) in place of methanol in the titration vessel.

Acceptance criteria: NMT 0.5%

• **OTHER REQUIREMENTS:** It meets the requirements for [Topical Aerosols \(603\)](#), in the sections *Pressure Test*, *Minimum Fill*, and *Leakage Test*.

ADDITIONAL REQUIREMENTS

• **PACKAGING AND STORAGE:** Preserve in pressurized containers, and avoid exposure to excessive heat.

• **USP REFERENCE STANDARDS (11).**

[USP Bacitracin Zinc RS](#)

[USP Polymyxin B Sulfate RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
BACITRACIN AND POLYMYXIN B SULFATE TOPICAL AEROSOL	Ying Han Associate Science & Standards Liaison	BIO42020 Biologics Monographs 4 - Antibiotics

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 28(4)

Current DocID: GUID-5BB4D572-99C6-4C70-B5F9-52106952F5C4_1_en-US

DOI: https://doi.org/10.31003/USPNF_M6885_01_01

DOI ref: [52eqq](#)

OFFICIAL