Status: Currently Official on 17-Feb-2025
Official Date: Official as of 01-Aug-2017
Document Type: USP Monographs
DocId: GUID-6644E114-1841-4D48-9C9E-3FBE20AC3DD7_1_en-US
DOI: https://doi.org/10.31003/USPNF_M8102_01_01
DOI Ref: bai3n

© 2025 USPC
Do not distribute

Benzocaine, Butamben, and Tetracaine Hydrochloride Topical Aerosol

DEFINITION

Benzocaine, Butamben, and Tetracaine Hydrochloride Topical Aerosol is Benzocaine, Butamben, and Tetracaine Hydrochloride Topical Solution packaged in a pressurized container with a suitable inert propellant. It contains NLT 90.0% and NMT 110.0% of the labeled amount of benzocaine ($C_9H_{11}NO_2$), butamben ($C_{11}H_{15}NO_2$), and tetracaine hydrochloride ($C_{15}H_{24}N_2O_2 \cdot HCl$).

IDENTIFICATION

- **A.** The retention times of the major peaks of the *Sample solution* correspond to those of the *Standard solution*, as obtained in the *Assay*.
- **B.** The UV spectrum of the major peaks of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the *Assay*.

ASSAY

- **Procedure**

Solution A: 0.1% formic acid in water

Solution B: 0.1% formic acid in acetonitrile

Mobile phase: See *Table 1*.

Table 1

Time (min)	Solution A (%)	Solution B (%)
0	91	9
2.5	50	50
3.9	50	50
4	91	9
5	91	9

Diluent: Acetonitrile and water (10:90)

Standard stock solution A: 1750 µg/mL of USP Benzocaine RS prepared as follows. Transfer a suitable amount of USP Benzocaine RS to a suitable volumetric flask and dissolve in 10% of the total volume of acetonitrile. Dilute with water to volume.

Standard stock solution B: 250 µg/mL each of USP Butamben RS and USP Tetracaine Hydrochloride RS prepared as follows. Transfer a suitable amount of USP Butamben RS and USP Tetracaine Hydrochloride RS to a suitable volumetric flask and dissolve in 10% of the total volume of acetonitrile. Dilute with water to volume.

Standard solution: 175 µg/mL of USP Benzocaine RS from *Standard stock solution A* and 25 µg/mL each of USP Butamben RS and USP Tetracaine Hydrochloride RS from *Standard stock solution B* diluted in *Diluent*

Sample solution: Nominally 175 µg/mL of benzocaine and 25 µg/mL each of butamben and tetracaine hydrochloride, prepared as follows. Accurately weigh about 125 mg of the evaporated sample into a 100-mL volumetric flask. Dissolve in 50 mL of methanol and dilute with *Diluent* to volume.

Chromatographic system

(See *Chromatography ⟨621⟩, System Suitability*.)

Mode: LC

Detector: UV 300 nm. For *Identification B*, use a diode array detector in the range of 200–400 nm.

Column: 2.1-mm × 5-cm; 1.7-µm packing L1

Flow rate: 0.6 mL/min

Injection volume: 1 µL

System suitability

Sample: *Standard solution*

[Note—The relative retention times for benzocaine, tetracaine, and butamben are about 0.71, 0.74, and 1.0, respectively.]

Suitability requirements

OFFICIAL

Documents
Document Sharing Hub
Full document details: Benzocaine, Butamben, and Tetracaine Hydrochloride Topical Aerosol

Resolution: NLT 2 between benzocaine and tetracaine

Relative standard deviation: NMT 2.0% for each of the three analyte peaks

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of benzocaine ($C_9H_{11}NO_2$), butamben ($C_{11}H_{15}NO_2$), and tetracaine hydrochloride ($C_{15}H_{24}N_2O_2 \cdot HCl$) in the portion of Topical Aerosol taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response of the corresponding analyte from the *Sample solution*

r_S = peak response of the corresponding analyte from the *Standard solution*

C_S = concentration of the corresponding Reference Standard in the *Standard solution* (µg/mL)

C_U = nominal concentration of the corresponding analyte in the *Sample solution* (µg/mL)

Acceptance criteria: 90.0%–110.0%

PERFORMANCE TESTS

• **Minimum Fill ⟨755⟩:** Meets the requirements

IMPURITIES

• **Organic Impurities**

Mobile phase, Diluent, and **Chromatographic system:** Proceed as directed in the *Assay*.

System suitability solution: 10 µg/mL each of USP Benzocaine RS, USP Tetracaine Hydrochloride RS, USP Butamben RS, and USP Ethyl 4-Nitrobenzoate RS in *Diluent*

Standard solution: 3.4 µg/mL each of USP Benzocaine RS and USP Ethyl 4-Nitrobenzoate RS and 1 µg/mL of USP Tetracaine Hydrochloride RS in *Diluent*

Sample solution: Nominally 1.68 mg/mL of benzocaine, 0.24 mg/mL of butamben, and 0.24 mg/mL of tetracaine prepared as follows. Accurately weigh about 600 mg of evaporated sample into a 50-mL volumetric flask, dissolve with 25 mL of methanol, and dilute with *Diluent* to volume. [Note—Sonication for about 1 min may be necessary.]

System suitability

Samples: *System suitability solution* and *Standard solution*

[Note—See *Table 2* for relative retention times.]

Suitability requirements

Resolution: NLT 2 between butamben and ethyl 4-nitrobenzoate; NLT 2 between benzocaine and tetracaine, *System suitability solution*

Relative standard deviation: NMT 5.0% for each of the analyte peaks, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of 4-aminobenzoic acid in the portion of Topical Aerosol taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (1/F) \times 100$$

r_U = peak response of 4-aminobenzoic acid from the *Sample solution*

r_S = peak response of benzocaine from the *Standard solution*

C_S = concentration of USP Benzocaine RS in the *Standard solution* (µg/mL)

C_U = nominal concentration of benzocaine in the *Sample solution* (µg/mL)

F = relative response factor (see *Table 2*)

Calculate the percentage of ethyl 4-nitrobenzoate in the portion of Topical Aerosol taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response of ethyl 4-nitrobenzoate from the *Sample solution*

r_S = peak response of ethyl 4-nitrobenzoate from the *Standard solution*

C_S = concentration of USP Ethyl 4-Nitrobenzoate RS in the *Standard solution* (µg/mL)

C_U = nominal concentration of benzocaine in the *Sample solution* (µg/mL)

OFFICIAL

AVA Documents Document Sharing Hub

Full document details: Benzocaine, Butamben, and Tetracaine Hydrochloride Topical Aerosol