


Status: Currently Official on 14-Feb-2025
 Official Date: Official as of 01-May-2020
 Document Type: USP Monographs
 DocId: GUID-0A286B6E-2103-4B1D-AA3A-8FD46DC280BA_4_en-US
 DOI: https://doi.org/10.31003/USPNF_M10790_04_01
 DOI Ref: qbu1n

© 2025 USPC
 Do not distribute

Butalbital



$C_{11}H_{16}N_2O_3$ 224.26
 2,4,6(1*H*,3*H*,5*H*)-Pyrimidinetrione, 5-(2-methylpropyl)-5-(2-propenyl)-;
 5-Allyl-5-isobutylbarbituric acid CAS RN®: 77-26-9; UNII: KHS0AZ4JVK.

DEFINITION

Butalbital contains NLT 98.0% and NMT 102.0% of butalbital ($C_{11}H_{16}N_2O_3$), calculated on the dried basis.

IDENTIFICATION

Change to read:

- **A.** [▲SPECTROSCOPIC IDENTIFICATION TESTS \(197\)](#), [Infrared Spectroscopy: 197K](#)▲ (CN 1-MAY-2020)
- **B.** The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.

ASSAY

• PROCEDURE

Buffer: 1.0 mL of phosphoric acid diluted with water to 1 L

Mobile phase: Acetonitrile and *Buffer* (25:75)

System suitability solution: 0.1 mg/mL each of [USP Butalbital RS](#) and [USP Salicylic Acid RS](#) in *Mobile phase*. Sonication may be used to aid in dissolution.

Standard solution: 0.1 mg/mL of [USP Butalbital RS](#) in *Mobile phase*. Sonication may be used to aid in dissolution.

Sample solution: 0.1 mg/mL of Butalbital in *Mobile phase*. Sonication may be used to aid in dissolution.

Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

Mode: LC

Detector: UV 214 nm

Column: 4.6-mm × 10-cm; 5-μm packing L1

Column temperature: 30°

Flow rate: 1 mL/min

Injection volume: 20 μL

System suitability

Samples: *System suitability solution* and *Standard solution*

[NOTE—The relative retention times of salicylic acid and butalbital are 0.86 and 1.0, respectively.]

Suitability requirements

Resolution: NLT 3.0 between salicylic acid and butalbital, *System suitability solution*

Tailing factor: NMT 1.5, *Standard solution*

Relative standard deviation: NMT 1.0%, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of butalbital ($C_{11}H_{16}N_2O_3$) in the portion of Butalbital taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of [USP Butalbital RS](#) in the *Standard solution* (mg/mL)

C_U = concentration of Butalbital in the *Sample solution* (mg/mL)

Acceptance criteria: 98.0%–102.0% on the dried basis

IMPURITIES

- [RESIDUE ON IGNITION \(281\)](#): NMT 0.1%

• ORGANIC IMPURITIES

Buffer: 4.1 g/L of monobasic potassium phosphate adjusted with 1 N sodium hydroxide to a pH of 6.0

Mobile phase: Acetonitrile and *Buffer* (22:78)

System suitability solution: 10 µg/mL each of [USP Butalbital RS](#) and [USP Butabarbital RS](#) in *Mobile phase*. Sonication may be used to aid in dissolution.

Sample solution: 1 mg/mL of Butalbital in *Mobile phase*. Sonication may be used to aid in dissolution.

Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

Mode: LC

Detector: UV 214 nm

Column: 4.6-mm × 15-cm; 5-µm packing L78

Column temperature: 30°

Flow rate: 1 mL/min

Injection volume: 20 µL

System suitability

Sample: *System suitability solution*

[NOTE—The relative retention times of butabarbital and butalbital are 0.83 and 1.0, respectively.]

Suitability requirements

Resolution: NLT 2.0 between butabarbital and butalbital

Tailing factor: NMT 1.5 for butalbital

Relative standard deviation: NMT 5.0% for butalbital

Analysis

Sample: *Sample solution*

Calculate the percentage of each impurity in the portion of Butalbital taken:

$$\text{Result} = (r_U/r_T) \times 100$$

r_U = peak response of each impurity from the *Sample solution*

r_T = sum of the peak responses from the *Sample solution*

Acceptance criteria

Any individual unspecified impurity: NMT 0.10%

Total impurities: NMT 1%

SPECIFIC TESTS

- [LOSS ON DRYING \(731\)](#)

Analysis: Dry under vacuum at room temperature to constant weight.

Acceptance criteria: NMT 0.2%

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in well-closed containers.

- [USP REFERENCE STANDARDS \(11\)](#)

[USP Butabarbital RS](#)

[USP Butalbital RS](#)

[USP Salicylic Acid RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
BUTALBITAL	Documentary Standards Support	SM42020 Small Molecules 4

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 38(4)