

Status: Currently Official on 14-Feb-2025
 Official Date: Official as of 01-Dec-2021
 Document Type: USP Monographs
 DocId: GUID-28975B79-FF9B-4B52-A781-564186C32A71_2_en-US
 DOI: https://doi.org/10.31003/USPNF_M11314_02_01
 DOI Ref: vg0ye

© 2025 USPC
 Do not distribute

Add the following:

^Calcipotriene Cream

DEFINITION

Calcipotriene Cream contains NLT 90.0% and NMT 110.0% of the labeled amount of calcipotriene ($C_{27}H_{40}O_3$).

IDENTIFICATION

- **A.** The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.
- **B.** The UV spectrum of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.

ASSAY

• PROCEDURE

Protect solutions containing calcipotriene from light and air. Prepare the solutions containing calcipotriene NMT 1 h before use.

Solution A: 0.132 g/mL of [monobasic ammonium phosphate](#) in [water](#)

Mobile phase: [Methanol](#) and [water](#) (70:30)

Diluent: [Methanol](#), *Solution A*, and [water](#) (70: 0.3: 29.7)

Standard stock solution: 0.1 mg/mL of [USP Calcipotriene RS](#) in *Diluent*. Sonicate if necessary.

Standard solution: 2.0 µg/mL of [USP Calcipotriene RS](#) prepared as follows. Transfer 5 mL of *Standard stock solution* into a 250-mL volumetric flask, add 50 mL of [tetrahydrofuran](#), and dilute with *Diluent* to volume.

Sample stock solution: Nominally equivalent to 0.01 mg/mL of calcipotriene in [tetrahydrofuran](#) prepared as follows. Transfer Cream nominally equivalent to 0.25 mg of calcipotriene to a 25-mL volumetric flask. Add 10 mL of [tetrahydrofuran](#) and sonicate for 20 min with intermittent shaking. Cool to room temperature and dilute with [tetrahydrofuran](#) to volume.

Sample solution: Nominally equivalent to 2.0 µg/mL of calcipotriene prepared as follows. Transfer 5 mL of *Sample stock solution* into a suitable container. Add 20 mL of *Diluent*, mix, and sonicate for 15 min. Pass through a Teflon filter of 0.45-µm pore size and discard the first few milliliters of the filtrate.

Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

Mode: LC

Detector: UV 264 nm. For *Identification B*, use a diode array detector in the range of 190–400 nm.

Column: 4.6-mm × 15-cm; 3-µm packing [L1](#)

Flow rate: 1.0 mL/min

Injection volume: 50 µL

System suitability

Sample: *Standard solution*

Suitability requirements

Tailing factor: NMT 2.0

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of calcipotriene ($C_{27}H_{40}O_3$) in the portion of Cream taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response of calcipotriene from the *Sample solution*

r_S = peak response of calcipotriene from the *Standard solution*

C_S = concentration of [USP Calcipotriene RS](#) in the *Standard solution* (mg/mL)

C_U = nominal concentration of calcipotriene in the *Sample solution* (mg/mL)

Acceptance criteria: 90.0%–110.0%

PERFORMANCE TESTS

- [MINIMUM FILL \(755\)](#): Meets the requirements

IMPURITIES**• ORGANIC IMPURITIES**

Protect solutions containing calcipotriene from light and air. Prepare the *Standard solution* and the *Sample solution* NMT 1 h before use.

Prepare the *System suitability solution* daily.

Mobile phase: See [Table 1](#).

Table 1

Time (min)	Water (%)	Methanol (%)
0	35	65
45	35	65
55	25	75
60	15	85
70	15	85
75	35	65
85	35	65

Diluent: Prepare as directed in the Assay.

System suitability stock solution: 0.125 mg/mL of [USP Calcipotriene RS](#) and 2.0 µg/mL of [USP Calcipotriene Related Compound C RS](#) in *Diluent*

System suitability solution: 0.0125 mg/mL of [USP Calcipotriene RS](#) and 0.2 µg/mL of [USP Calcipotriene Related Compound C RS](#) prepared as follows. Transfer 1 mL of *System suitability stock solution* to a 10-mL volumetric flask. Add 1 mL of [tetrahydrofuran](#) and dilute with *Diluent* to volume.

Standard stock solution: 62.5 µg/mL of [USP Calcipotriene RS](#) prepared as follows. Transfer an appropriate amount of [USP Calcipotriene RS](#) to a suitable volumetric flask, add *Diluent* equivalent to 50% of the final volume, and sonicate to dissolve. Further add [tetrahydrofuran](#) equivalent to 25% of the final volume and dilute with *Diluent* to volume.

Standard solution: 0.125 µg/mL of [USP Calcipotriene RS](#) in *Diluent*, from *Standard stock solution*

Sensitivity solution: 0.0125 µg/mL of [USP Calcipotriene RS](#) in *Diluent*, from *Standard solution*

Sample solution: Nominally 0.0125 mg/mL of calcipotriene prepared as follows. Transfer Cream equivalent to 0.25 mg of calcipotriene to a 20-mL volumetric flask and add 5 mL of [tetrahydrofuran](#). Sonicate for 15 min with intermittent shaking. Cool to room temperature and dilute with *Diluent* to volume. Place the volumetric flask in a beaker containing ice-cold water for 2–3 min, and pass the solution through a 0.45-µm Teflon filter. Discard the first few milliliters of the filtrate.

Chromatographic system

(See [Chromatography \(621\)](#), *System Suitability*.)

Mode: LC

Detector: UV 264 nm

Column: 4.6-mm × 15-cm; 3-µm packing [L1](#)

Column temperature: 30°

Flow rate: 1.0 mL/min

Injection volume: 100 µL

System suitability

Samples: *System suitability solution*, *Standard solution*, and *Sensitivity solution*

Suitability requirements

Resolution: NLT 1.2 between calcipotriene related compound C and calcipotriene, *System suitability solution*

Relative standard deviation: NMT 5.0%, *Standard solution*

Signal-to-noise ratio: NLT 10, *Sensitivity solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of each individual impurity in the portion of Cream taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (1/F) \times 100$$

r_U = peak response of each individual impurity from the *Sample solution*