

RECOMMENDATION ITU-R P.371-8

CHOICE OF INDICES FOR LONG-TERM
IONOSPHERIC PREDICTIONS

(1963-1970-1974-1978-1982-1986-1990-1995-1999)

The ITU Radiocommunication Assembly,

recommends

- 1 that the 12-month running mean sunspot number R_{12} , or alternatively the 12-month running mean value of Φ , the 2800 MHz solar radio noise flux (i.e. Φ_{12}), be adopted as the preferred index to be used for predicting monthly median values of foF2 and M(3000)F2 over all time-scales: substantially equivalent results should be obtainable by the use of either of these indices;
- 2 that Φ_{12} be adopted as the preferred index to be used for predicting monthly median values of foE and foF1 over all time-scales;
- 3 that predicted values of these indices should be determined by means of the modified McNish-Lincoln procedure (see Annex 1) using latest available measured monthly index values for the present solar cycle and the average of past cycle values for future cycles;
- 4 that where propagation predictions require simultaneous use of values of different ionospheric characteristics, the same index may be adopted for all such characteristics with little loss of accuracy;
- 5 that caution be shown in the use of the recommended indices at high magnetic latitudes, where the resulting ionospheric predictions may not be sufficiently accurate.

ANNEX 1

1 Introduction

The concept of indices for long-term ionospheric predictions relies on the assumption that the important characteristics of the ionosphere, such as the critical frequencies of the various layers and the MUF factor M(3000)F2, depend in a systematic way on certain measurable quantities concerned with solar radiation. It should however be noted that the correlation between these indices and the actual ionospheric characteristics does not necessarily imply a causal relationship, but rather an indication of associated phenomena. Changes of solar activity in general contain three components:

- a fairly regular component with a period of about 11 years, which represents the well known cycle of solar activity;
- a component that has a quasi-period of about a year or a little less; and
- erratic fluctuations with periods of less than a month.

2 Sunspot numbers

For studies of the main component of the solar cycle, the 12-month running mean sunspot number R_{12} is used because the resultant smoothing considerably reduces complicated rapidly-varying components, but does not obscure the slowly-varying component.

The definition of R_{12} is:

$$R_{12} = \frac{1}{12} \left[\sum_{k=n-5}^{n+5} R_k + \frac{1}{2} (R_{n+6} + R_{n-6}) \right] \quad (1)$$

in which R_k is the mean of the daily sunspot numbers for a single month k , and R_{12} is the smoothed index for the month represented by $k = n$.

The two main disadvantages in the use of R_{12} are:

- the most recent available value is necessarily centred on a month at least 6 months earlier than the present time;
- it cannot be used to predict the shorter-term variation in solar activity.

Nonetheless, R_{12} appears to be the most useful parameter for long-term studies and predictions concerning the F2 layer.

3 Index Φ

Consistent and reasonably long series of observations of the solar radio noise flux at about 10 cm wavelength have been made by Canadian, Japanese and other laboratories. The monthly mean, Φ , of the daily values from Canada, expressed in units of $10^{-22} \text{ Wm}^{-2} \text{ Hz}^{-1}$, should be regarded as the reference data for this index. Φ is more closely correlated with E-layer critical frequency than are noise flux values at other wavelengths. As the solar flux observations are only available from 1947, the sunspot numbers remain one of the longest series of observations of a natural phenomenon. Therefore, the continued collection and recording of sunspot observations is encouraged.

4 Other indices

Over past years, a substantial number of different indices have been considered to try to represent long-term changes of the different ionospheric characteristics, but of these the ITU-R recommends the indices R_{12} and Φ_{12} for ionospheric predictions.

5 Correlation between Φ_{12} and R_{12}

The recommended relationship between R_{12} and Φ_{12} , also indicated in Fig. 1, is:

$$\Phi_{12} = 63.7 + 0.728 R_{12} + 8.9 \times 10^{-4} R_{12}^2 \quad (2)$$

6 The prediction of indices

There is as yet no method whereby it is possible to predict accurately indices for the next sunspot cycle, or, more generally, for a cycle which has not yet begun. Indices that have been calculated by using harmonic analysis, or by using empirical and statistical laws applied to observations over some earlier and even recent cycles, have not proved useful in predicting those for a new cycle. After a sunspot minimum has been observed, future development of the cycle can be extrapolated to a certain extent, although the deviations have been observed to be rather extreme.