

Status: Currently Official on 15-Feb-2025
Official Date: Official as of 01-Dec-2017
Document Type: USP Monographs
DocId: GUID-6A2EC5DB-D2F6-4693-BD4F-65D87BC13A30_1_en-US
DOI: https://doi.org/10.31003/USPNF_M9797_01_01
DOI Ref: wv6dn

© 2025 USPC
Do not distribute

Leucovorin Calcium Compounded Oral Suspension

DEFINITION
Leucovorin Calcium Compounded Oral Suspension contains NLT 90.0% and NMT 110.0% of the labeled amount of leucovorin ($C_{20}H_{23}N_7O_7$).
Prepare Leucovorin Calcium Compounded Oral Suspension containing 5 mg/mL of leucovorin as follows (see [Pharmaceutical Compounding—Nonsterile Preparations \(795\)](#)).

Leucovorin Calcium tablets ^a equivalent to	500 mg of leucovorin
Sodium Hydroxide (1 N)	To adjust pH to 7.1–7.6
Syrup, a sufficient quantity to make	100 mL

^a Leucovorin calcium 25-mg tablets, Teva Pharmaceuticals, Sellersville, PA.

Place the *Leucovorin Calcium tablets* into a suitable container. Wet the tablets with a small amount of *Syrup* and triturate to make a smooth paste. Add the *Syrup* to make the contents pourable. Transfer contents stepwise and quantitatively to a calibrated container using the *Syrup*. Adjust with *Sodium Hydroxide (1 N)* to a pH of 7.1–7.6. Add sufficient *Syrup* to bring to final volume. Shake to mix well. [NOTE—pH may decrease to 6.1 after bringing to final volume with *Syrup* without affecting the stability of the preparation.]

ASSAY

• **PROCEDURE**

Solution A: Methanol and 5 mM tetrabutylammonium phosphate (20:80). Adjust with tetrabutylammonium hydroxide to a pH of 6.6. [NOTE—Tetrabutylammonium phosphate appearing wet should not be used as it may coelute with leucovorin.]
Mobile phase: See [Table 1](#).

Table 1

Time (min)	Methanol (%)	Solution A (%)
0	0	100
20	10	90
20.1	0	100
30	0	100

Diluent: Methanol and water (20:80)
Standard solution: 0.05 mg/mL of leucovorin prepared from [USP Leucovorin Calcium RS](#) and *Diluent*. Vortex and sonicate until dissolved.
Sample solution: Transfer 1.0 mL of Oral Suspension to a 100-mL volumetric flask, and rinse the pipette with about 2 mL of *Diluent*. Dilute with *Diluent* to volume.
Chromatographic system
(See [Chromatography \(621\), System Suitability](#).)
Mode: LC
Detector: UV 290 nm
Column: 4.6-mm × 15-cm; 2.7-μm packing L7
Column temperature: 55°



Flow rate: 0.75 mL/min

Injection volume: 10 µL

System suitability

Sample: *Standard solution*

[NOTE—The retention time for leucovorin is about 20.3 min.]

Suitability requirements

Tailing factor: NMT 2.0

Relative standard deviation: NMT 2.0% for replicate injections

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of leucovorin ($C_{20}H_{23}N_7O_7$) in the portion of Oral Suspension taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of leucovorin from [USP Leucovorin Calcium RS](#) in the *Standard solution* (mg/mL)

C_U = nominal concentration of leucovorin in the *Sample solution* (mg/mL)

Acceptance criteria: 90.0%–110.0%

SPECIFIC TESTS

- **pH** (791): 6.1–7.1

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Package in tight, light-resistant plastic containers. Store in a refrigerator or at controlled room temperature.
- **BEYOND-USE DATE:** NMT 90 days after the date on which it was compounded when stored in a refrigerator; NMT 30 days after the date on which it was compounded when stored at controlled room temperature
- **LABELING:** Label it to be well shaken before use, and to state the *Beyond-Use Date*.
- **USP REFERENCE STANDARDS** (11).
[USP Leucovorin Calcium RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
LEUCOVORIN CALCIUM COMPOUNDED ORAL SUSPENSION	Brian Serumaga Science Program Manager	CMP2020 Compounding 2020
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	CMP2020 Compounding 2020

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 41(2)

Current DocID: GUID-6A2EC5DB-D2F6-4693-BD4F-65D87BC13A30_1_en-US

DOI: https://doi.org/10.31003/USPNF_M9797_01_01

DOI ref: [wv6dn](#)