Status: Currently Official on 16-Feb-2025
Official Date: Official as of 01-Aug-2017
Document Type: USP Monographs
DocId: GUID-BF5407E0-C675-4CFD-A3BB-21DB0C789397_1_en-US
DOI: https://doi.org/10.31003/USPNF_M56131_01_01
DOI Ref: e1mnz

© 2025 USPC
Do not distribute

Neomycin and Polymyxin B Sulfates, Bacitracin, and Hydrocortisone Acetate Ophthalmic Ointment

DEFINITION

Neomycin and Polymyxin B Sulfates, Bacitracin, and Hydrocortisone Acetate Ophthalmic Ointment contains the equivalent of NLT 90.0% and NMT 140.0% of the labeled amounts of neomycin, polymyxin B, and bacitracin, and NLT 90.0% and NMT 110.0% of the labeled amount of hydrocortisone acetate in a suitable ointment base.

IDENTIFICATION

• **A. Thin-Layer Chromatographic Identification Test ⟨201BNP⟩:** Meets the requirements

• **B.** The retention time of the hydrocortisone acetate peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the *Assay* for *Hydrocortisone Acetate*.

ASSAY

• **Neomycin**

(See *Antibiotics—Microbial Assays ⟨81⟩*.)

Sample solution: Shake a portion of Ophthalmic Ointment in a separator with 50 mL of ether. Extract with four 20-mL portions of *Buffer B.3*. Combine the aqueous extracts, and dilute with *Buffer B.3* to a suitable volume.

Analysis: Proceed as directed in the chapter. Dilute the *Sample solution* with *Buffer B.3* to obtain a *Test Dilution* having a neomycin concentration that is nominally equivalent to the median level of the standard.

Acceptance criteria: 90.0%–140.0%

• **Polymyxin B**

(See *Antibiotics—Microbial Assays ⟨81⟩*.)

Sample solution: Shake a portion of Ophthalmic Ointment with 50 mL of ether in a separator. Extract with four 25-mL portions of *Buffer B.6*. Combine the aqueous extracts, and dilute with *Buffer B.6* to a suitable volume.

Analysis: Proceed as directed in the chapter. Dilute the *Sample solution* with *Buffer B.6* to obtain a *Test Dilution* having a concentration that is nominally equivalent to the median level of the standard (10 polymyxin B units/mL). Add to each *Test Dilution* of the standard a quantity of USP Neomycin Sulfate RS, dissolved in *Buffer B.6*, to obtain the same concentration of neomycin as in the *Test Dilution* of the sample.

Acceptance criteria: 90.0%–140.0%

• **Bacitracin**

(See *Antibiotics—Microbial Assays ⟨81⟩*.)

Sample solution: Shake a portion of Ophthalmic Ointment with 50 mL of ether in a separator. Extract with four 20-mL portions of *Buffer B.1*. Combine the buffer extracts, and dilute with *Buffer B.1* to a suitable volume.

Analysis: Proceed as directed in the chapter. Add sufficient 0.01 N hydrochloric acid to a portion of the *Sample solution* so that the amount of hydrochloric acid in the *Test Dilution* is the same as in the median level of the standard. Dilute with *Buffer B.1* to obtain a *Test Dilution* having a bacitracin concentration that is nominally equivalent to the median level of the standard.

Acceptance criteria: 90.0%–140.0%

• **Hydrocortisone Acetate**

Mobile phase: Butyl chloride, water-saturated butyl chloride, tetrahydrofuran, methanol, and glacial acetic acid (475:475:70:35:30)

Standard solution: 0.10 mg/mL of USP Hydrocortisone Acetate RS in water-saturated chloroform

Sample solution: Nominally 0.10 mg/mL of hydrocortisone acetate from Ophthalmic Ointment prepared as follows. Transfer a portion of Ophthalmic Ointment containing nominally 2.5 mg of hydrocortisone acetate to a closable container. Add 25.0 mL of water-saturated chloroform and about 10 glass beads. Securely close the container, and shake vigorously for approximately 15 min. Centrifuge, and use the clear, lower chloroform layer.

Chromatographic system

(See *Chromatography ⟨621⟩, System Suitability*.)

Mode: LC

OFFICIAL

Documents
Document Sharing Hub

Full document details: Neomycin and Polymyxin B Sulfates, Bacitracin, and Hydrocortisone Acetate Ophthalmic Ointment

Detector: UV 254 nm

Column: 3.9-mm × 30-cm; 10-µm packing L3

System suitability

Sample: *Standard solution*

Suitability requirements

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of hydrocortisone acetate ($C_{23}H_{32}O_6$) in the portion of Ophthalmic Ointment taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of USP Hydrocortisone Acetate RS in the *Standard solution* (mg/mL)

C_U = nominal concentration of hydrocortisone acetate in the *Sample solution* (mg/mL)

Acceptance criteria: 90.0%–110.0%

SPECIFIC TESTS

• **Sterility Tests ⟨71⟩:** Meets the requirements

• **Other Requirements:** It meets the requirements for *Particulate and Foreign Matter* and *Container Contents* in *Ophthalmic Products—Quality Tests ⟨771⟩, Drug Product Quality, Universal Tests, Particulate and Foreign Matter* and *Container Contents*.

ADDITIONAL REQUIREMENTS

• **Packaging and Storage:** Preserve in collapsible ophthalmic ointment tubes.

• **USP Reference Standards ⟨11⟩**

USP Bacitracin Zinc RS
USP Hydrocortisone Acetate RS
USP Neomycin Sulfate RS
USP Polymyxin B Sulfate RS

Auxiliary Information - Please check for your question in the FAQs before contacting USP.

Topic/Question	Contact	Expert Committee
NEOMYCIN AND POLYMYXIN B SULFATES, BACITRACIN, AND HYDROCORTISONE ACETATE OPHTHALMIC OINTMENT	Julie Zhang Associate Science & Standards Liaison	BIO42020 Biologics Monographs 4 - Antibiotics
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	BIO42020 Biologics Monographs 4 - Antibiotics

Chromatographic Database Information: Chromatographic Database

Most Recently Appeared In:
Pharmacopeial Forum: Volume No. PF 39(5)

Current DocID: GUID-BF5407E0-C675-4CFD-A3BB-21DB0C789397_1_en-US

DOI: https://doi.org/10.31003/USPNF_M56131_01_01

DOI ref: e1mnz

OFFICIAL

Documents
Document Sharing Hub

Full document details: Neomycin and Polymyxin B Sulfates, Bacitracin, and Hydrocortisone Acetate Ophthalmic Ointment