Status: Currently Official on 16-Feb-2025
Official Date: Official as of 01-Aug-2017
Document Type: USP Monographs
DocId: GUID-70592DA8-D767-4817-B58E-6E762B3CF9FF_1_en-US
DOI: https://doi.org/10.31003/USPNF_M56220_01_01
DOI Ref: z3g89

© 2025 USPC
Do not distribute

Neomycin and Polymyxin B Sulfates, Bacitracin Zinc, and Hydrocortisone Acetate Ophthalmic Ointment

DEFINITION

Neomycin and Polymyxin B Sulfates, Bacitracin Zinc, and Hydrocortisone Acetate Ophthalmic Ointment is a sterile ointment containing Neomycin Sulfate, Polymyxin B Sulfate, Bacitracin Zinc, and Hydrocortisone Acetate. It contains the equivalent of NLT 90.0% and NMT 140.0% of the labeled amounts of neomycin, polymyxin B, and bacitracin, and NLT 90.0% and NMT 110.0% of the labeled amount of hydrocortisone acetate ($C_{23}H_{32}O_6$).

IDENTIFICATION

• **A. Thin-Layer Chromatographic Identification Test ⟨201BNP⟩:** Meets the requirements

• **B.** The retention time of the hydrocortisone acetate peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the *Assay* for *Hydrocortisone Acetate*.

ASSAY

• **Neomycin**

(See *Antibiotics—Microbial Assays ⟨81⟩*.)

Sample solution: Shake a portion of Ophthalmic Ointment in a separator with 50 mL of ether. Extract with four 20-mL portions of *Buffer B.3*. Combine the aqueous extracts, and dilute with *Buffer B.3* to a suitable volume.

Analysis: Proceed as directed in the chapter. Dilute the *Sample solution* with *Buffer B.3* to obtain a *Test Dilution* having a neomycin concentration that is nominally equivalent to the median level of the standard.

Acceptance criteria: 90.0%–140.0%

• **Polymyxin B**

(See *Antibiotics—Microbial Assays ⟨81⟩*.)

Sample solution: Shake a portion of Ophthalmic Ointment with 50 mL of ether in a separator. Extract with four 25-mL portions of *Buffer B.6*. Combine the aqueous extracts, and dilute with *Buffer B.6* to a suitable volume.

Analysis: Proceed as directed in the chapter. Dilute the *Sample solution* with *Buffer B.6* to obtain a *Test Dilution* having a concentration that is nominally equivalent to the median level of the standard (10 polymyxin B Units/mL). Add to each *Test Dilution* of the standard a quantity of USP Neomycin Sulfate RS, dissolved in *Buffer B.6*, to obtain the same concentration of neomycin as in the *Test Dilution* of the sample.

Acceptance criteria: 90.0%–140.0%

• **Bacitracin**

(See *Antibiotics—Microbial Assays ⟨81⟩*.)

Sample solution: Shake a portion of Ophthalmic Ointment with 50 mL of ether in a separator. Extract with four 25-mL portions of 0.01 N hydrochloric acid. Combine the acid extracts, and dilute with 0.01 N hydrochloric acid to a suitable volume.

Analysis: Proceed as directed in the chapter. Dilute the *Sample solution* with *Buffer B.1* to obtain a *Test Dilution* having a concentration that is nominally equivalent to the median level of the standard (1.0 bacitracin Unit/mL). If the *Sample solution* has a concentration of less than 100 bacitracin Units/mL, add hydrochloric acid to each *Test Dilution* of the standard to obtain the same concentration of hydrochloric acid as in the *Test Dilution* of the sample.

Acceptance criteria: 90.0%–140.0%

• **Hydrocortisone Acetate**

Mobile phase: Butyl chloride, water-saturated butyl chloride, tetrahydrofuran, methanol, and glacial acetic acid (475:475:70:35:30)

Standard solution: 0.10 mg/mL of USP Hydrocortisone Acetate RS in water-saturated chloroform

Sample solution: Nominally 0.10 mg/mL of hydrocortisone acetate from Ophthalmic Ointment prepared as follows. Transfer a portion of Ophthalmic Ointment containing nominally 2.5 mg of hydrocortisone acetate to a closable container. Add 25.0 mL of water-saturated chloroform and about 10 glass beads. Securely close the container, and shake vigorously for approximately 15 min. Centrifuge, and use the clear, lower chloroform layer.

OFFICIAL
AVA Documents
Document Sharing Hub
Full document details: Neomycin and Polymyxin B Sulfates, Bacitracin Zinc, and Hydrocortisone Acetate Ophthalmic Ointment

Chromatographic system

(See *Chromatography ⟨621⟩, System Suitability*.)

Mode: LC

Detector: UV 254 nm

Column: 3.9-mm × 30-cm; 10-µm packing L3

System suitability

Sample: *Standard solution*

Suitability requirements

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of hydrocortisone acetate ($C_{23}H_{32}O_6$) in the portion of Ophthalmic Ointment taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of USP Hydrocortisone Acetate RS in the *Standard solution* (mg/mL)

C_U = nominal concentration of hydrocortisone acetate in the *Sample solution* (mg/mL)

Acceptance criteria: 90.0%–110.0%

SPECIFIC TESTS

• **Sterility Tests ⟨71⟩:** Meets the requirements

• **Other Requirements:** It meets the requirements for *Particulate and Foreign Matter* and *Container Contents* in *Ophthalmic Products—Quality Tests ⟨771⟩, Drug Product Quality, Universal Tests, Particulate and Foreign Matter* and *Container Contents*.

ADDITIONAL REQUIREMENTS

• **Packaging and Storage:** Preserve in collapsible ophthalmic ointment tubes. Store at controlled room temperature.

• **USP Reference Standards ⟨11⟩**

USP Bacitracin Zinc RS
USP Hydrocortisone Acetate RS
USP Neomycin Sulfate RS
USP Polymyxin B Sulfate RS

Auxiliary Information - Please check for your question in the FAQs before contacting USP.

Topic/Question	Contact	Expert Committee
NEOMYCIN AND POLYMYXIN B SULFATES, BACITRACIN ZINC, AND HYDROCORTISONE ACETATE OPHTHALMIC OINTMENT	Julie Zhang Associate Science & Standards Liaison	BIO42020 Biologics Monographs 4 - Antibiotics
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	BIO42020 Biologics Monographs 4 - Antibiotics

Chromatographic Database Information: Chromatographic Database

Most Recently Appeared In:
Pharmacopeial Forum: Volume No. PF 39(5)

Current DocID: GUID-70592DA8-D767-4817-B58E-6E762B3CF9FF_1_en-US

DOI: https://doi.org/10.31003/USPNF_M56220_01_01

DOI ref: z3g89

OFFICIAL

Documents
Document Sharing Hub

Full document details: Neomycin and Polymyxin B Sulfates, Bacitracin Zinc, and Hydrocortisone Acetate Ophthalmic Ointment